



# Respiratory Pathogens


*Is it a Cold, RSV, Influenza or COVID-19?*




Rhinovirus (Common Cold)



Respiratory Syncytial Virus (RSV)



Influenza (Flu Virus)



COVID-19

	Rhinovirus (Common Cold)	Respiratory Syncytial Virus (RSV)	Influenza (Flu Virus)	COVID -19
Causative agent (pathogen)	Many different viruses including rhinoviruses (most common), parainfluenza and seasonal coronaviruses.	Respiratory Syncytial Virus (RSV)	Influenza viruses types A & B	SARS-CoV-2
How it spreads	Respiratory droplets and direct contact with the infected person, their respiratory secretions or stool ("poop"), or with contaminated surfaces, followed by touching eyes, nose or mouth.	Respiratory droplets and direct contact with the infected person or their respiratory secretions, or with contaminated surfaces followed by touching eyes, nose or mouth.	Person-to-person by way of large respiratory droplets and direct contact with their respiratory secretions, or contaminated surfaces, followed by touching eyes, nose or mouth.	Primarily through breathing in small droplets or particles that contain the virus, or when they land on eyes, nose or mouth through splashes, sprays-like coughs or sneezes.  Contact with their respiratory secretions or contaminated surfaces, followed by touching eyes, nose or mouth.
Contagious period	24 hours before symptom onset to 5 days afterward	3-8 days but can be as long as 4 weeks.	From 1 day before symptom onset to 5-7 days afterward.	From 2 days before symptom onset to -10 days afterward. This includes asymptomatic persons.
Incubation period (time to symptom onset after an exposure)	-2 days	4-6 days	1-4 days	2-14 days
Symptoms	Symptom onset is gradual. Common symptoms include: <ul style="list-style-type: none"> <li>▶ Sneezing</li> <li>▶ Mild to moderate chest discomfort, cough</li> <li>▶ Stuffy nose</li> <li>▶ Sore throat</li> </ul>	Symptoms appear in stages and not all at once and include: <ul style="list-style-type: none"> <li>▶ Runny nose</li> <li>▶ Decreased appetite</li> <li>▶ Coughing</li> <li>▶ Sneezing</li> <li>▶ Fever</li> <li>▶ Wheezing</li> </ul> In very young infants, the only symptoms might be: <ul style="list-style-type: none"> <li>▶ Irritability</li> <li>▶ Decreased activity</li> <li>▶ Difficulty breathing</li> </ul>	Symptom onset is abrupt. Common symptoms include: <ul style="list-style-type: none"> <li>▶ Fever, chills</li> <li>▶ Aches</li> <li>▶ Fatigue, weakness</li> <li>▶ Chest discomfort, cough</li> <li>▶ Headache</li> </ul>	Symptoms range from none to severe and include: <ul style="list-style-type: none"> <li>▶ Fever, chills</li> <li>▶ Cough</li> <li>▶ Shortness of breath or difficulty breathing</li> <li>▶ Fatigue</li> <li>▶ Muscle aches</li> <li>▶ Headache</li> <li>▶ New loss of taste or smell</li> <li>▶ Congestion, runny nose</li> <li>▶ Nausea, vomiting, diarrhea</li> </ul>

	Rhinovirus (Common Cold)	Respiratory Syncytial Virus (RSV)	Influenza (Flu Virus)	COVID -19
<b>Survival on surfaces</b>	2 hours to 7 days (rhinovirus)	7-8 hours	2 hours to several weeks	4 hours to 8 days
<b>Vaccine preventable</b>	No	No	Yes	Yes
<b>Infection prevention</b>	▶ Frequent hand hygiene	▶ Frequent hand hygiene	▶ Get vaccinated annually.	▶ Get vaccinated
	▶ Avoid touching eyes, nose, mouth with unclean hands	▶ Avoid touching eyes, nose, mouth with unclean hands	▶ Frequent hand hygiene	▶ Wear a mask
	▶ Cover coughs, sneezes	▶ Cover cough, sneezes	▶ Avoid touching eyes, nose, mouth with unclean hands	▶ Social distance (6 feet from others)
	▶ Avoid contact with people who are sick	▶ Avoid contact people who are sick	▶ Cover coughs, sneezes	▶ Frequent hand hygiene
	▶ Clean & disinfect surfaces and objects that may be contaminated with the virus	▶ Clean & disinfect surfaces and objects that may be contaminated with the virus	▶ Avoid contact with people who are sick	▶ Avoid touching eyes, nose, mouth with unclean hands
	▶ Stay home when ill	▶ Stay home when ill	▶ Clean & disinfect surfaces and objects that may be contaminated with the virus	▶ Cover coughs and sneezes
			▶ Stay home when ill	▶ Avoid crowds and poorly ventilated spaces
				▶ Avoid contact with people who are sick
			▶ Clean & disinfect high touch surfaces daily	
			▶ Stay home when ill	

**References:**

Centers for Disease Control & Prevention. Influenza (Flu) [Internet]. [Cited 2021 Nov 18]. Available from <https://www.cdc.gov/flu/index.htm>

Centers for Disease Control & Prevention. Common Colds: Protect Yourself and Others [Internet]. [Cited 2021 Nov 18]. Available from <https://www.cdc.gov/features/rhinoviruses/index.html>

Heymann DL. Control of Communicable Diseases Manual. 19th edition. Washington, DC: American Public Health Association; 2008. 516-517 p.

Wissmann JE et al. Persistence of Pathogens in Inanimate Surfaces: A Narrative Review. Microorganisms. 2021 Feb 9;9(2):343.

Centers for Disease Control and Prevention. Respiratory Syncytial Virus (RSV) [Internet]. [Cited 2021 Nov 18. Available from <https://www.cdc.gov/rsv/index.html>

Centers for Disease Control and Prevention. About COVID-19 [Internet]. [Cited 2021 Nov 18]. Available from <https://www.cdc.gov/coronavirus/2019-ncov/your-health/about-covid-19.html>

Compiled by Doe Kley, RN, MPH, CIC, T-CHEST, Sr. Infection Preventionist, Clorox Healthcare

